



Lac La Biche County
welcoming by nature.

2021 Water Quality Report Beaver Lake

Lac La Biche County
Alberta, Canada

Prepared by:
Lac La Biche County
Box 1679
Lac La Biche, AB T0A 2C0

Contributions: Randi Dupras and Nolan Ekeberg

Executive Summary

Beaver lake is a large and attractive recreational lake located in Lac La Biche County, Alberta (“County”) and is popular for a variety of recreational activities. However, there is a concern that declining water quality in the lake is limiting the opportunities of recreational activities like swimming, boating, and fishing.

The County follows a regular program to monitor water quality of lakes located within its jurisdiction. The water sampling events were conducted during the early spring and summer of 2021. The data collected includes water temperature, pH, specific conductivity, and dissolved oxygen which was collected in-situ through a multi-probe and Kemmerer sampling device. Analytical data of nitrogenous compounds, heavy metals, and other inorganic parameters was provided from ALS laboratory.

Collected water samples were analyzed by ALS laboratory. The laboratory results obtained were compared to the CCME’s Canadian Environmental Quality Guidelines for Protection of Aquatic Life and Protection of Agricultural Water, and Alberta Environment and Parks’ Environmental Quality Guidelines for Alberta Surface Waters 2018.

Trophic State Index (TSI) is a classification system designed to rate lakes based on the amount of biological activity they sustain. The concentrations of nutrients (nitrogen and phosphorous) are the primary determinants of TSI. Increased concentrations of nutrients tend to result in increased plant growth, followed by an increase in subsequent trophic level. Nurnberg (1996) used parameters including Secchi depth, chlorophyll, total nitrogen, and total phosphorus concentrations in lake waters to determine the trophic state of the lakes, which is provided as Table 1 in Appendix A. TSI is a useful tool for evaluation and management of lake health and setting objectives including sport and recreational activities related to the lake. Trophic classification of Beaver Lake based on Secchi depth and nutrients is presented in Table 2 in Appendix A.

For the purpose of this report, the parameters used to determine the trophic state will only include Secchi depth, total nitrogen and total phosphorus. Chlorophyll will not be used to determine the trophic state. Chlorophyll is a green pigment present in all green plants and is responsible for the absorption of light to provide energy for photosynthesis. It is associated with algae growth in a waterbody and affects the trophic status of a lake. Chlorophyll concentration is measured as part of the County’s monitoring program. However, the measurement can be an underestimate of algae biomass when blue-green algae are present. It is also difficult to have consistent measurements of Chlorophyll as there can be large variances in concentrations due to anomalies such as temperature and weather conditions. Therefore, it is difficult to report Chlorophyll concentrations and make recommendations based on the results. Based on this information, Chlorophyll is not reported in this document.

There are four classes of trophic states which include: Oligotrophic which would be the highest quality of water with low productivity, nutrients, and algae; Mesotrophic which is fair quality water with some productivity, nutrients and algae; Eutrophic which is relatively poor-quality water with high productivity, nutrients and algae; and Hypereutrophic which is the poorest quality water with excessive productivity, nutrients, and algae.

Beaver Lake would be considered Eutrophic based on the average of the three water parameters Secchi depth, total nitrogen, and total phosphorus. The trophic status would be Mesotrophic based on Secchi depth, Hypereutrophic based on total nitrogen, and Eutrophic based on total phosphorus.

Results and Discussion

In 2021, Secchi depths in Beaver Lake were measured on March 9, June 22, July 26, and August 19, 2021. The average seasonal Secchi depth was observed to be 2.7 m, which is slightly higher with historical results. The low average Secchi depth means that Beaver lake water has poor transparency due to suspended materials. Based on the Secchi depths and in accordance with the classification provided in Table 1 (Appendix A), Beaver Lake is classified as Mesotrophic (some productivity, nutrients, and algae growth).

Sampling events in 2021 showed an average summer water temperature of 18.75 °C. Uniform temperature profiles were observed during the summer as there were no significant variation in temperatures with depth. Based on the data provided, thermal stratification was not observed in any of the summer sampling events between March 9 and August 19, 2021.

Dissolved oxygen data collected in 2021 shows that the average dissolved oxygen levels ranged from 0.3 mg/L to 10.1 mg/L. These concentrations were within proximity to the regulatory criteria for dissolved oxygen in cold water lakes for early life stages (9.5 mg/L) and for all other life stages (6.5 mg/L).

In 2021, two types of lake water samples for analyses of nutrients were collected from Beaver Lake; composite samples and Kemmerer samples (obtained from different depths using a Kemmerer device). These samples were analyzed for total nitrogen and total phosphorous.

Total nitrogen concentrations in the composite samples collected from the lake in 2021 had an average of 1.29 mg/L of total nitrogen, while the Kemmerer samples collected had an average of 1.58 mg/L of total nitrogen; both of which exceeded the applicable regulatory guidelines and were consistent with historical results. Total nitrogen concentrations from both sampling methods classify Beaver Lake as Hypereutrophic (excessive productivity, nutrients, and algae growth).

Total phosphorus concentrations in the composite samples collected during the summer of 2021 had an average of 0.031 mg/L of total phosphorus, while the Kemmerer samples collected had an average of 0.031 mg/L; both of which do not exceed the applicable regulatory guidelines of 0.05 mg/L and were consistent with historical results. Total phosphorus concentrations from both sampling methods classify Beaver Lake as Eutrophic (high productivity, nutrients, and algae growth).

The average N:P ratios for composite and Kemmerer sampling events were 42:1 and 50:1 which is higher than the Redfield Ratio of 16:1. Therefore, the total phosphorus concentrations are considered low enough for phosphorus to be considered the main nutrient limiting growth in Beaver Lake.

Routine water chemistry showed that Beaver Lake has an average pH of 7.87 in 2021 which is consistent with historical results.

Concentrations of metals analyzed from the composite and Kemmerer samples taken at a depth of 6 m were generally below detection limits and/or below the applicable regulatory guidelines.

Beaver Lake would be considered Eutrophic based on the average of the three water parameters Secchi depth, total nitrogen, and total phosphorus. The trophic status would be Mesotrophic based on Secchi depth, Hypereutrophic based on total nitrogen, and Eutrophic based on total phosphorus.

Recommendations:

It is recommended that Lac La Biche County continues to monitor the water quality of Beaver Lake on a regular basis. Continuous monitoring will help the County to determine how the lake management strategies and policies such as the Watershed Management Plan and Riparian Setback Matrix Model are impacting the lake water quality, and what the net effect on human and environmental health is.

Monitoring and sampling should continue to be conducted under a strategic plan and in a uniform manner to ensure that results produced are meaningful and are useful for establishing a correlation with the past results. This may include sampling at same period of the year each time, recording the same parameters critical to lake health, obtaining samples from the same depths, and implementing a quality assurance program for reliability of analytical results.

Nutrient loading is the main source of eutrophication in Beaver Lake which is degrading the water quality; leading to algae growth, foul smells and a reduction in water recreation. Therefore, action must be taken to slow down the eutrophication process and improve water quality. Best management practices would include education of the public on appropriate land use including watershed protection and waste and recycling management; restoration and protection of riparian areas (water buffers); and strengthening laws and regulations governing land use such as municipal sewer hookups and protection of environmental reserves.

Lac La Biche County updated the Lac La Biche Watershed Management Plan, which was adopted by Council in May 2021. This plan will include specific action items based on the recommendations that are formulated while drafting the plan. Although Beaver Lake is not within the Lac La Biche watershed, the recommended action items may still apply.

Table of Contents

Executive Summary1
Results and Discussion.....2
Recommendations..... 2

1. INTRODUCTION.....6

2. WATER QUALITY SAMPLING PROGRAM6
2.1 Water Quality Parameters.....7

3. REGULATORY FRAMEWORK.....8

4. SAMPLING ANALYSIS AND MONITORING RESULTS8
4.1 Secchi Depth8
4.2 Temperature9
4.3 Dissolved Oxygen10
4.4 Nutrients11
4.5 Routine Water Chemistry15
4.6 Metals.....15
4.7 Alberta Lake Management Society Sampling.....16

5. HISTORICAL TREND ANALYSIS16
5.1 Secchi Depth16
5.2 Total Nitrogen17
5.3 Total Phosphorus.....18

6. DISCUSSION18

7. RECOMMENDATIONS19

8. REFERENCES.....20

APPENDIX A22

List of Figures

- Figure 1: Location map of Beaver Lake
- Figure 2: Secchi depths measured in Beaver Lake - 2021
- Figure 3: Temperature in Beaver Lake - 2021
- Figure 4: Dissolved oxygen in Beaver Lake - 2021
- Figure 5: Total nitrogen from composite samples of Beaver Lake - 2021
- Figure 6: Total nitrogen from Kemmerer samples of Beaver Lake - 2021
- Figure 7: Total phosphorous from composite samples of Beaver Lake - 2021
- Figure 8: Total phosphorus from Kemmerer samples of Beaver Lake - 2021
- Figure 9: Historical trend for Secchi depth in Beaver Lake
- Figure 10: Historical trend for total nitrogen in Beaver Lake
- Figure 11: Historical trend for total phosphorus in Beaver Lake
- Figure 12: Historical trend for water levels in Beaver Lake

List of Tables

- Table 1: Trophic status classification based on lake water parameters (Nurnberg 1996)
- Table 2: Trophic status of Beaver Lake based on lake water parameters - 2021
- Table 3: Average N:P ratios for Beaver Lake water samples in - 2021
- Table 4: Routine water chemistry parameters in Beaver Lake composite samples - 2021
- Table 5: Total dissolved metals in Kemmerer samples from Beaver Lake - 2021
- Table 6: Historical data of routine water chemistry in Beaver Lake
- Table 7: Historical data of total dissolved metals in Beaver Lake

Abbreviations/Acronyms Used

- CCME: Canadian Council of Ministers of the Environment
- County: Lac La Biche County
- EQGASW-AGW: Environmental Quality Guidelines for Alberta Surface Waters 2018 for protection of Agricultural Water
- EQGASW-FAL: Environmental Quality Guidelines for Alberta Surface Waters 2018 for protection of Fresh Water Aquatic Life
- QA/QC: Quality Assurance and Quality Control
- Total N: Total Nitrogen
- Total P: Total Phosphorous
- TSI: Trophic State Index

1. INTRODUCTION

Beaver lake is a large recreational lake popular for boating and fishing. It is located approximately 210 km northeast of the city of Edmonton. The closest population center is the hamlet of Lac La Biche which is located 5 km to the northwest of Beaver Lake. A location map of Beaver Lake is provided in Figure 1.




Figure 1: Location map of Beaver Lake

Beaver lake has an irregular shaped surface area of 33.1 Km² and is comprised of two distinct basins. The two basins are connected by a shallow and a narrow channel. The maximum depth of the lake is 15.2 m in the northwest basin and 10.7 m in the southeast basin. Beaver Lake is the headwater of the Beaver River and lies in the Beaver River Drainage Basin. The drainage basin for Beaver Lake is 290 km², which is roughly nine times the surface area of the lake itself. Beaver Lake receives water from two other lakes within the watershed; Elinor Lake which drains into the Southeast basin and Lac La Croix which drains into the Northwest Basin. In years of significantly high-water levels, Beaver Lake also receives water from Roseland and Normandeau Lakes. The lake's outlet creek, which is located on the west side of the north basin, flows into Outlet Lake, which further drains into the Beaver River and eventually into the Churchill River.

The main agricultural activities are livestock production, production of pasture and forage crops such as hay, oats and barley. A large number of natural gas and oil wells are found within the County.

The main sport fish species found in the lake include: Walleye (*Sander vitreus*), Yellow Perch (*Perca flavescens*), Northern Pike (*Esox lucius*) and Lake Whitefish (*Coregonus clupeaformis*).

2. WATER QUALITY SAMPLING PROGRAM

Lac La Biche County has sampled Beaver Lake from 2003 to 2005, 2010 to 2015, and 2017 to 2021. Beaver Lake is sampled for various parameters using different techniques. Vertical profiles were taken using a multi-probe testing different depths (zones) of the lake for dissolved oxygen, pH, conductivity, and

temperature. Composite samples are taken from 10 different locations throughout the lake, while Kemmerer sampling is used for discrete depth sampling; both the composite and Kemmerer samples are tested for nutrients such as, phosphorus, nitrogen, ammonia, nitrates, nitrites, and metals. Beaver Lake sampling program for 2021 was completed as follows:

- a) Secchi Depths were measured on June 22, July 26, and August 19, 2021.
- b) Composite samples from the Beaver Lake were collected on June 22, July 26, and August 19, 2021. Lake water samples were analyzed for nutrients, metals, and basic water chemistry parameters by ALS laboratories.
- c) Kemmerer water samples using the Kemmerer device were collected on March 9 and August 19, 2021 from depths of 3 m, 6 m, and 9 m; and were analyzed for nutrients, metals, and basic water chemistry parameters by ALS laboratories.
- d) Lake profiles were recorded to a maximum depth of 10 m using a multi-probe June 22, July 26, and August 19, 2021.

2.1 Water Quality Parameters

Water samples collected for each of the sampling locations were analyzed for a variety of parameters used to characterize the chemical composition of the lake and further identify any potential concerns. The water quality parameters measured and analyzed during the 2021 program along with a brief description of each parameter and reason for monitoring are provided in the table below:

Parameters Affecting Lake Water Quality

Water Quality Parameter	Description and Reason for Measuring
Secchi Depth	Secchi depth is a measure of the transparency of water and trophic state of a lake. A Secchi disk is generally a disk of 20 cm diameter with alternating black and white quadrants. It is lowered into the lake water until it can no longer be seen. This depth of disappearance is called the Secchi depth. A low Secchi depth (<4 m) is characteristic of a mesotrophic to hypereutrophic lake with turbid water. Whereas a high Secchi depth (>4 m) is characteristic of an oligotrophic lake with clear water.
Dissolved Oxygen	Dissolved oxygen is required by aquatic plants and animals for respiration. Survival of aquatic life such as fish, generally depends on an adequate amount of dissolved oxygen for respiration. As dissolved oxygen levels in the water drop below 5.0 mg/L, aquatic life is subjected to stress. Oxygen levels that consistently remain below 1-2 mg/L can result in the loss of large populations of fish.
Temperature	Temperature of water affects different physical, biological and chemical characteristics of a lake and determines the behavior of many parameters responsible for water quality. The solubility of oxygen and other gases decrease as temperature increases. An increase in water temperature decreases the concentration of dissolved oxygen required for the survival of aquatic organisms.

Nutrients	Total nitrogen (N) and phosphorus (P) are principal nutrients in lake water and are representative of all forms of N and P present in the water. There are various sources of N and P both natural and anthropogenic. These nutrients are a major cause of eutrophication, decreasing dissolved oxygen concentrations and are detrimental to lake water quality.
Metals	Metals enter the lake waters through natural (geological) and anthropogenic point and non-point sources. Certain metals such as lead and mercury, are toxic to aquatic life and can bio-accumulate in the tissues and organs of aquatic organisms, becoming a part of the food chain. This may lead to loss of aquatic life and further affect human health.

3. REGULATORY FRAMEWORK

The protection of water quality in Canadian lakes is a federal, provincial, and territorial responsibility. Therefore, lake waters in Alberta are regulated by federal and provincial guidelines and fall under the jurisdiction of Canadian Council of Ministers of the Environment (CCME), Alberta Environment and Parks (AEP), and Health Canada.

The regulatory criteria selection for lake waters in Alberta are subjected to CCME’s Canadian Environmental Quality Guidelines (CEQG) and AEP’s Environmental Quality Guidelines for Alberta Surface Waters 2018 (EQGASW). Protection of lake water is covered under CCME’s CEQG and AEP’s EQGASW chapters of water quality guidelines for Protection of Aquatic Life, Protection of Agricultural Water, and protection of Recreation and Aesthetics. In addition, Health Canada’s Guidelines for Canadian Recreational Water Quality for protection of lake waters have also been considered.

The analytical and monitoring results obtained for this report were compared to the above-mentioned regulations and are hereinafter referred to as regulatory guidelines or regulatory criteria.

4. SAMPLING ANALYSIS AND MONITORING RESULTS

4.1 Secchi Depth

The Secchi disk is a common method used to measure water clarity. Water clarity of a lake can be influenced by the amount of suspended materials such as phytoplankton, zooplankton, pollen, sediments and dissolved compounds. The Secchi depth multiplied by 2 provides us with the euphotic depth of the lake. The euphotic depth is the maximum depth to which light can penetrate within a lake to facilitate growth.

In 2021, Secchi depths in Beaver Lake were measured on June 22, July 26, and August 19, 2021. The average seasonal Secchi depth was observed to be 2.7 m. A maximum Secchi depth of 4.5 m was recorded on June 22, 2021 while a minimum Secchi depth of 1.5 m was recorded on August 19, 2021. Overall, a decreasing temporal trend was observed in Secchi depth (Figure 2).

The low average Secchi depth of 2.7 m means that the lake water has poor transparency due to suspended materials. Based on the Secchi depths and in accordance with the classification provided in Table 1 (Appendix A), Beaver Lake is classified as Mesotrophic (some productivity, nutrients, and algae growth).




Figure 2: Secchi depths measured in Beaver Lake - 2021

4.2 Temperature

Water temperature in a lake determines the behavior of many parameters responsible for water quality. Thermal stratification occurs within a lake with a distinct difference in temperature between the surface water (epilimnion layer) and the deeper water (hypolimnion layer) separated by a thermocline. The thermocline is identified when the water changes by more than one degree Celsius per meter. Under winter conditions, ice covers the surface water, and a thermocline is formed with the colder water at the surface and the warmer water at the bottom of the lake. Lakes without thermal stratification mix from top to bottom and this mixing allows oxygen to distribute throughout the water column preventing hypolimnetic anoxia (lack of oxygen). In the summertime, warmer surface water can facilitate cyanobacteria blooms at the lake surface (Wetzel, R. 2001).

The temperatures in Beaver Lake were recorded to a maximum depth of 12 m. A minimum average temperature of 2.70 °C was observed on March 9, 2021 with a spatial increasing trend with depth, starting from 0.867 °C at 1 m and 4.404 °C at 12 m depth.

Sampling events in 2021 showed an average summer water temperature of 18.75 °C. Uniform temperature profiles were observed during the summer as there were no significant variation in temperatures with depth. Based on the data provided, thermal stratification was not observed in any of the summer sampling events between June 6 and August 19, 2021. Results of temperatures observed at varying depths at the sampling dates are illustrated in Figure 3.

Similar trends in temperature profiles have been reported for Beaver Lake previously (ALMS 2010). Historical sampling also suggests that Beaver Lake is likely polymictic, indicating that the water from various depths of the lake mixes several times per year.




Figure 3: Temperature in Beaver Lake – 2021

4.3 Dissolved Oxygen

Dissolved oxygen is the amount of gaseous oxygen dissolved in the water and is necessary for respiration and survival of aquatic life (e.g., fish, invertebrates, bacteria, and underwater plants). Dissolved oxygen is also needed for the decomposition of organic matter in the lakes. Oxygen enters the lake water by direct absorption from the atmosphere through rapid movement of water or as a product of plant photosynthesis. Therefore, the epilimnion zone (shallow layer of water) is relatively richer in oxygen than the hypolimnion zone (deeper layer of water) which is low in oxygen due to consumption by respiration.

There are several conditions necessary for fish survival in a lake including adequate water temperatures and available dissolved oxygen for respiration. The regulatory guidelines for dissolved oxygen in cold water lakes are 9.5 mg/L for early life stages and 6.5 mg/L for all other life stages (CCME, 1999). If dissolved oxygen levels are too low, fish will move to other depths in the water column, often where temperatures are conducive to sustain aquatic life.

The amount of dissolved oxygen in lakes usually decreases under winter ice-cover primarily due to respiration by organisms (particularly bacteria) and decomposition of organic matter. In shallow lakes, oxygen depletion can proceed rapidly under ice during the winter. If dissolved oxygen drops below 3.0 mg/L during the winter, many fish and invertebrate species will not survive.

Dissolved oxygen levels in Beaver Lake were recorded to a maximum depth of 11 m using a multi-probe on March 9, June 22, July 26, and August 19, 2021. A maximum concentration of 10.07 mg/L of dissolved oxygen was observed on July 26, 2021 at a depth of 1 m depth, which declined gradually to 6.36 mg/L at a depth of 11 m.

A decrease in dissolved oxygen was noted in the winter measurements with a decreasing gradual spatial trend with depth; the decreasing gradual spatial trend was minimal in the summer measurements. The lowest level of dissolved oxygen was recorded on March 9, 2021 (0.00 mg/L) at a 11 m depth (lakebed).

Dissolved oxygen data collected in 2021 shows that average dissolved oxygen levels ranged from 4.23 mg/L on March 9, 2021 to 8.40 mg/L on August 19, 2021. These concentrations were in proximity to the

regulatory criteria for dissolved oxygen in cold water lakes for early life stages (9.5 mg/L) and for all other life stages (6.5 mg/L).




Figure 4: Dissolved oxygen in Beaver Lake – 2021

4.4 Nutrients

Excessive levels of nitrogen and phosphorus are found in many lakes across Alberta leading to excessive growth of algae and aquatic plants. Decay of aquatic vegetation causes oxygen depletion in the water column and contributes to eutrophication. Consequently, the decreased levels of oxygen can suffocate fish and other aquatic organisms. High nutrient conditions foster algal blooms and can result in the proliferation of toxin-producing blue-green algae (e.g., cyanobacteria). The input of nutrients into aquatic systems can occur naturally, but large amounts of nutrients typically originate from indirect, non-point anthropogenic sources, including improperly treated sewage, residential use of fertilizers and agricultural operations.

In 2021, two types of lake water samples for analyses of nutrients were collected from Beaver Lake; composite samples and Kemmerer samples (obtained from different depths using a Kemmerer device). These samples were analyzed for total nitrogen and total phosphorus.

Total Nitrogen

Total nitrogen is an essential nutrient for plants and animals; however, excessive amounts of nitrogen in lake water may lead to low levels of dissolved oxygen and negatively affect water quality and health of aquatic life within the lake. Nitrogen concentrations in the water are typically measured in three forms: ammonia, nitrates, and nitrites. Total nitrogen is the sum of total Kjeldahl nitrogen (ammonia, organic and reduced nitrogen), nitrate and nitrite. Nitrogen levels in lakes are also affected by atmospheric deposition, which refers to nitrogen in the air being deposited into the water system. Nitrogen oxides (NOx) are added to atmosphere due to the burning of fossil fuels, so emissions from motor vehicles and industrial facilities can also affect nitrogen levels in aquatic environments.

Composite Samples

Composite lake water samples for analyses of total nitrogen were collected on June 22, July 26, and August 19, 2021. The total nitrogen concentrations ranged from 1.09 mg/L to 1.44 mg/L during 2021. The analytical results are displayed in Figure 5.

Nitrogen concentrations in the composite samples collected from the lake in 2021 had an average of 1.29 mg/L of total nitrogen which exceeds the applicable regulatory guidelines. The average total nitrogen indicates that Beaver Lake is Hypereutrophic (excessive productivity, nutrients, and algae growth) based on total nitrogen from composite samples.




Figure 5: Total nitrogen from composite samples of Beaver Lake – 2021

Kemmerer Sampling

Kemmerer water samples are collected from different depths of the lake by using a Kemmerer device which makes it possible to obtain a sample of water from specific depths. Kemmerer samples were collected on March 9 and August 19, 2021 from 3 m, 6 m, and 9 m and were analyzed for total nitrogen by ALS laboratories. Results of total nitrogen in Kemmerer samples collected from Beaver Lake on two different dates are illustrated in Figure 6. On March 9, 2021 a total nitrogen concentration of 1.9 mg/L was recorded at a depth of 6 m, which increased to a concentration of 2.33 mg/L at a depth of 9 m. On August 19, 2021 a total nitrogen concentration of 1.29 mg/L was recorded at a depth of 3 m, which decreased to a concentration of 1.12 mg/L at a depth of 9 m.

The average concentrations of total nitrogen in samples collected during March 9 and August 19, 2021 sampling events was 1.58 mg/L which exceeds the applicable regulatory guidelines. The results from the Kemmerer sampling resulted in the same trophic state classification as the composite samples which is Hypereutrophic (excessive productivity, nutrients, and algae growth) based on total nitrogen.




Figure 6: Total nitrogen from Kemmerer samples of Beaver Lake – 2021

Total Phosphorus

Increased phosphorus concentrations are the largest cause of degradation in water quality within lakes, causing ‘dead zones’, toxic algal blooms, a loss of biodiversity and increased health risks for plants, animals and humans that encounter polluted lake waters. Run-off from agriculture, human sewage and industrial practices results in increased phosphorus concentrations in lake water and lakebed sediments (Wetzel, 2001). Long-term monitoring activities following the control of phosphorus sources to lakes indicates that plants and animals do not recover from the effects of excessive phosphorus for several years.

Composite Sampling

Composite lake water samples for total phosphorus were collected on June 22, July 26, and August 19, 2021 from Beaver Lake. A minimum total phosphorus concentration of <0.020 mg/L was measured in a sample obtained on June 6, 2021. The highest concentration of phosphorus was 0.036 mg/L which was observed on August 19, 2021. The analytical results are illustrated below in Figure 7.

Total phosphorus concentrations in the samples collected during the summer of 2021 had an average of 0.031 mg/L of total phosphorus which does not exceed the applicable regulatory guidelines of 0.05 mg/L. This average total phosphorus concentration classifies Beaver Lake as Eutrophic (high productivity, nutrients, and algae growth) based on total phosphorus from composite samples.




Figure 7: Total phosphorous from composite samples of Beaver Lake – 2021

Kemmerer Sampling

Kemmerer water samples using a Kemmerer sampling device were collected on March 9 and August 19, 2021 from depths of 3 m, 6 m, and 9 m. Both samples were analyzed for total phosphorous by ALS laboratories. The sample from the 3m depth on March 9, 2021 did not have results provided from the water quality analysis, so there are no values at this depth. Total phosphorus concentrations in lake water samples collected on March 9, 2021 had an average concentration of 0.026 mg/L and was slightly lower than the average concentration of samples collected on August 19, 2021 which was 0.035 mg/L. The laboratory results are presented below in Figure 8.

Total phosphorus concentrations from the Kemmerer sampling in 2021 did not exceed the regulatory criteria. The results from the Kemmerer sampling resulted in the same trophic state classification as the composite samples which is Eutrophic (high productivity, nutrients, and algae growth) based on total phosphorus.




Figure 8: Total phosphorus from Kemmerer samples of Beaver Lake - 2021

N:P Ratio

The Redfield Ratio describes the optimal balance of total nitrogen to total phosphorous for aquatic plant growth, which is an optimal value of 16:1 (Teubner and Dokulil 2002). If the ratio is lower than 16:1, phosphorus is no longer considered a limiting nutrient and aquatic vegetation and cyanobacteria can use the dissolved and atmospheric nitrogen for growth by using the high amounts of phosphorus available in lake waters. If the ratio is higher than 16:1, it indicates that the phosphorus concentrations are occurring at levels much less than nitrogen and hence limit the growth within lakes.

The average N:P ratios for composite and Kemmerer sampling events in Beaver Lake were 42:1 and 50:1 respectively, which is higher than the Redfield Ratio of 16:1. Therefore, the total phosphorus concentrations are considered low enough for phosphorous to be considered the main nutrient limiting growth in the Beaver Lake.

4.5 Routine Water Chemistry

Results of routine water chemistry of composite and Kemmerer samples collected from Beaver Lake are presented in Table 4.

The average measured pH of 7.87 in 2021 was consistent with historical results. The pH of water determines the solubility and biological availability of chemical constituents such as nutrients and heavy metals. The ability of a lake to neutralize these hydrogen ions is referred to as a buffering capacity. Any lake with a total alkalinity of more than 100 mg/L is considered to have high buffering capacity (Mitchell and Prepas 1990). The pH in Beaver Lake is likely buffered against change by its high alkalinity. The high alkalinity in Alberta lakes is derived from the rich calcareous glacial till over which the lakes have formed.

4.6 Metals

Metals enter the water naturally through the weathering of rocks and soil. These metals are generally non-toxic and in low concentrations. However, metals can also come from a wide variety of anthropogenic and non-point pollution sources including runoff from urban areas, wastewater discharge, improperly managed sewage treatment, industrial activities, and agricultural runoff. The analytical results of total dissolved

metals in the Kemmerer and composite water samples collected from Beaver Lake are presented in Table 5.

Concentrations of all metals analyzed from the composite and Kemmerer samples taken at a depth of 9 m were generally below detection limits and/or below the applicable regulatory guidelines.

4.7 Alberta Lake Management Society Sampling

During the summer of 2021, Alberta Lake Management Society (ALMS) worked with Lac La Biche County to complete watermilfoil sampling along with quagga & zebra mussel sampling, and spiny water flea (aquatic invasive species) in Beaver Lake. During the last week of August, Lac La Biche County collected a watermilfoil sample from Beaver Lake. The sample was sent to Alberta Plant Health Laboratory and was analyzed to determine if the specimen was native Northern Watermilfoil (*Myriophyllum sibiricum*) or the invasive Eurasian Watermilfoil (*Myriophyllum spicatum*). The sample from Beaver Lake was determined to be native Northern Watermilfoil.

5. HISTORICAL TREND ANALYSIS

The objective of the historical trend analysis is to provide an overview of water quality conditions in a lake with time and to evaluate the impact of watershed management practices on lake water quality.

Three parameters are significant in trend analyses for lake water quality: Secchi depth, total nitrogen and total phosphorus; all of which are also used for trophic classification of lakes. A graph of Beaver Lake water level has also been included for review.

5.1 Secchi Depth

Historical data indicates that the Secchi depth in Beaver Lake was always less than the standard Oligotrophic (low productivity, nutrients, and algae growth) standard for Secchi depth (4.0 m) except for 2010 when Secchi depth (4.2 m) was slightly higher than the standard (Figure 9). A temporal decreasing trend in Secchi depth was observed for Beaver Lake following 2012, until 2019, and 2020, when the average Secchi depth slightly improved. However, the overall trend for Secchi depth shows a gradual deterioration of water quality. The low average Secchi depth means that the lake water has poor transparency due to suspended materials. However, the Secchi depth readings may not provide an exact measure of the water transparency due to various errors such as time of the day, sun's glare on the water, and eyesight of the observer.




Figure 9: Historical trend for Secchi depth in Beaver Lake

5.2 Total Nitrogen

Historical data shows that total nitrogen concentrations in Beaver Lake were historically higher than the regulatory guideline of 1.0 mg/L. A gradual increasing trend in total nitrogen has been observed from 1986 to 2021 and a sharp increase was measured in 2013. Total nitrogen concentrations have historically been classified as Hypereutrophic (excessive productivity, nutrients, and algae growth) as seen in Figure 10 below.




Figure 10: Historical trend for total nitrogen in Beaver Lake

5.3 Total Phosphorus

Historical data shows that total phosphorus concentrations in Beaver Lake were less than the regulatory guidelines (0.05 mg/L), except for 2012 (0.07 mg/L). An overall trend of increasing total phosphorus concentrations has been observed; however, the concentrations are below the regulatory criteria, aside from 2012 (Figure 11). Total phosphorus concentrations have historically been classified as Eutrophic (high productivity, nutrients, and algae growth).




Figure 11: Historical trend for total phosphorus in Beaver Lake

6. DISCUSSION

Trophic State Index (TSI) is a classification system designed to rate lakes based on the amount of biological activity they sustain. The concentrations of nutrients (nitrogen and phosphorous) are the primary determinants of TSI. Increased concentrations of nutrients tend to result in increased plant growth, followed by an increase in subsequent trophic level. Nurnberg (1996) used parameters including Secchi depth, chlorophyll, total nitrogen and total phosphorus concentrations in lake waters to determine the trophic state of the lakes, which is provided as Table 1 in Appendix A. TSI is a useful tool for evaluation and management of lake health and setting objectives including sport and recreational activities related to the lake. Trophic classification of Beaver Lake based on Secchi depth and nutrients is presented in Table 2.

For the purpose of this report, the parameters used to determine the trophic state will only include Secchi depth, total nitrogen and total phosphorus. Chlorophyll will not be used to determine the trophic state. Chlorophyll is a green pigment present in all green plants and is responsible for the absorption of light to provide energy for photosynthesis. It is associated with algae growth in a waterbody and affects the trophic status of a lake. Chlorophyll concentration is measured as part of the County’s monitoring program. However, the measurement can be an underestimate of algae biomass when blue-green algae are present. It is also difficult to have consistent measurements of Chlorophyll as there can be large variances in concentrations due to anomalies such as temperature and weather conditions such as precipitation and wind. Therefore, it is difficult to report Chlorophyll concentrations and make recommendations based on the results. Based on this information, Chlorophyll is not reported in this document.

There are four classes of trophic states which include: Oligotrophic which would be the highest quality of water with low productivity, nutrients, and algae; Mesotrophic which is fair quality water with some productivity, nutrients, and algae; Eutrophic which is relatively poor-quality water with high productivity, nutrients and algae; and Hypereutrophic which is the poorest quality water with excessive productivity, nutrients, and algae.

Beaver Lake would be considered Eutrophic based on the average of the three water parameters Secchi depth, total nitrogen, and total phosphorus. The trophic status would be Mesotrophic based on Secchi depth, Hypereutrophic based on total nitrogen, and Eutrophic based on total phosphorus.

7. RECOMMENDATIONS

It is recommended that Lac La Biche County continues to monitor the water quality of Beaver Lake on a regular basis. Continuous monitoring will help the County to determine how the lake management strategies and policies such as the Watershed Management Plan and Riparian Setback Matrix Model are impacting the lake water quality, and what the net effect on human and environmental health is.

Monitoring and sampling should continue to be conducted under a strategic plan and in a uniform manner to ensure that results produced are meaningful and are useful for establishing a correlation with the past results. This may include sampling at same period of the year each time, recording the same parameters critical to lake health, obtaining samples from the same depths, and implementing a quality assurance program for reliability of analytical results.

Nutrient loading is the main source of eutrophication in Beaver Lake which is degrading the water quality, leading to algae growth, foul smells and a reduction in water recreation. Therefore, action must be taken to slow down the eutrophication process and improve water quality. Best management practices would include education of the public on appropriate land use including watershed protection and waste and recycling management; restoration and protection of riparian areas (water buffers); and strengthening laws and regulations governing land use such as municipal sewer hookups and protection of environmental reserves.

Lac La Biche County updated the Lac La Biche Watershed Management Plan, which was adopted by Council in May 2021. This plan will include specific action items based on the recommendations that are formulated while drafting the plan. Although Beaver Lake is not within the Lac La Biche watershed, the recommended action items may still apply.

8. REFERENCES

1. Atlas of Alberta Lakes, 1990. <http://albertalakes.ualberta.ca/?page=home>, accessed September 22, 2018
2. Baby, J., J. S. RAJ, E. T. Biby, P. Sankarganesh, M.V. Jeevitha, S.U. Ajisha and S. S. Rajan, Toxic effect of heavy metals on aquatic environment. *Int. J. Biol. Chem. Sci.* 4(4): 939-952, 2010
3. Burns N. M. and, Nriagu, J. O., Forms of Iron and Manganese in Lake Erie Waters, *Journal of the Fisheries Research Board of Canada*, 1976, 33(3): 463-470, 2011
4. Canadian Council of Ministers of the Environment. 1999. Canadian water quality guidelines for the protection of aquatic life: Dissolved oxygen (freshwater). In: Canadian environmental quality guidelines, Canadian Council of Ministers of the Environment, Winnipeg, 1999
5. Canadian Council of Ministers of the Environment, Canadian Environmental Quality Guidelines, 2007
6. Government of Alberta, Alberta Guide to Sport Fishing Regulations, 2018
7. Government of Alberta, Environmental Quality Guidelines for Alberta Surface Waters, Alberta Environment and Parks, 2018
8. Elayse M. Hachich,* Marisa Di Bari, Ana Paula G. Christ, Cláudia C. Lamparelli, Solange S. Ramos, and Maria Inês Z. Sato; Comparison of thermotolerant coliforms and *Escherichia coli* densities in freshwater bodies, *Brazilian Journal of Microbiology*; 43(2): 675–681; 2012
9. Government of Alberta, Guide to the commercial fishing seasons, 2012
10. Government of Alberta, <http://aep.alberta.ca/fish-wildlife/default.aspx>, accessed September 22, 2018
11. Government of Alberta. (2019). *Alberta Safe Beach Protocol*. Created by Alberta Health, Public Health and Compliance. Retrieved from <https://open.alberta.ca/publications/9781460145395>
12. Government of Alberta, Trophic state of Alberta lakes based on average total chlorophyll, 2013. <https://open.alberta.ca/publications/trophic-state-of-alberta-lakes-based-on-average-chlorophyll-a-concentrations>, accessed on September 22, 2018
13. Government of Alberta, Trophic state of Alberta lakes based on average total phosphorus concentrations, 2013. <https://open.alberta.ca/publications/trophic-state-of-alberta-lakes-based-on-average-total-phosphorus-concentrations>, accessed September 22, 2018
14. Health Canada Guidelines for Canadian Recreational Water Quality, 2012
15. Lac La Biche County Office, Lac La Biche East and West, Water Sampling Report, 2016
16. Mitchell, P.A. and E.E. Prepas (eds.), Atlas of Alberta Lakes, University of Alberta Press. (detailed information on 100 Alberta lakes: author of introduction on Water Quality and six lake chapters, co-author on nine lake chapters) p.690, 1990
17. Nurnberg, G. 1996. Trophic state of clear and colored, soft- and hardwater lakes with special consideration of nutrients, anoxia, phytoplankton and fish. *Lake Reserv. Man.* 12(4): 432-447.
18. Schindler, D. W. et al, The cultural eutrophication of Lac la Biche, Alberta, Canada: a paleoecological study. *Can. J. Fish. Aquat. Sci.* 65: 2211–2223, 2008
19. Teubner, K. and M. T. Dukulil, Ecological stoichiometry of TN:TP:SRSi in freshwaters: nutrient ratios and seasonal shifts in phytoplankton assemblages. *Arch Hydrobiol.* 625-646, 2002
20. Thrane, J. E., D. O. Hessen, and T. Andersen 2014. The Absorption of Light in Lakes: Negative Impact of Dissolved Organic Carbon on Primary Productivity. *Ecosystems* 17: 1040–1052, 2014
21. Thurston, R. V., C. R. Rosemarie, and G. A. Vinogradov, 1981. Ammonia toxicity to fish; Effect of pH on the toxicity of the unionized ammonia species. *Environ. Sci. & Technol.* 15 (7): 837-840

22. Wade, T. J., Calderon, R.L., Brenner, K. P., Sams, E., Beach, M.J., Haugland, R., ... Dufour, A.P. High sensitivity of children to swimming-associated gastrointestinal illness: Results using a rapid assay of recreational water quality. *Epidemiology* 2008, 19 (3), 375-383.
23. Wetzel, R. G., *Limnology: Lake and River Ecosystems*, 3rd Edition; Elsevier Academic Press. 20, 2001

APPENDIX A

Table 1: Trophic status classification based on lake water parameters (Nurnberg 1996)

Trophic State	Total Phosphorus (mg/L)	Total Nitrogen (mg/L)	Secchi Depth (m)
Oligotrophic	<0.01	<0.35	>4
Mesotrophic	0.01 – 0.03	0.35 – 0.65	4 - 2
Eutrophic	0.03 – 0.10	0.65 – 1.20	2 - 1
Hypereutrophic	>0.1	>1.20	<1

Table 2: Trophic status of Beaver Lake based on lake water parameters – 2021

Trophic State	Secchi Depth	Total Nitrogen	Total Phosphorus
	(m)	------(mg/L) -----	
Oligotrophic	>4	<0.35	<0.01
Mesotrophic	4 – 2	0.35 – 0.65	0.01 – 0.03
Eutrophic	2 – 1	0.65 – 1.2	0.03 – 0.1
Hypereutrophic	<1	>1.2	>0.1
Beaver Lake 2021	2.7	1.29	0.031
Trophic State of Beaver Lake 2021	Mesotrophic	Hypereutrophic	Eutrophic
Trophic State of Beaver Lake 2020	Mesotrophic	Hypereutrophic	Eutrophic

Table 3: Average N:P ratios for Beaver Lake water samples in - 2021

Sampling Event	Total Nitrogen (mg/L)	Total Phosphorus (mg/L)	N:P
Composite Sampling	1.29	0.031	42:1
Kemmerer Sampling	1.58	0.031	50:1

Table 4: Routine water chemistry parameters in Beaver Lake composite samples – 2021

Date of Sampling	June 22, 2021	July 28, 2021	August 19, 2021
	mg/L		
pH	7.91	8.10	8.18
Temperature (°C)	17.59	19.39	19.28
Ammonia, Total (as N)	<0.050	<0.050	<0.050
Nitrate (as N)	<0.020	<0.020	<0.020
Nitrite (as N)	<0.010	<0.010	<0.010
Nitrate and Nitrite (as N)	<0.022	<0.022	<0.022

* Based on average pH and temperature of 7.87 and 18.75 °C of Beaver Lake in 2021

1: CCME C Guidelines, de-minimis criteria for Protection of Aquatic Life and Agricultural Water Uses

2 - Environmental Quality Guidelines for Alberta Surface Waters 2018

a: CCME Canadian Environmental Quality Guidelines for water for the Protection of Aquatic Life

b: CCME Guidelines for Protection of Agricultural Water Uses (Irrigation and Livestock pathways included)

Table 5: Total dissolved metals in Kemmerer samples from Beaver Lake – 2021

Date of Sampling	Kemmerer Sampling (9 m depth) March 09, 2021	Kemmerer Sampling (9 m depth) August 19, 2021	Criteria ¹	Criteria ²
Parameters	------(mg/L)-----			
Aluminum (Al)-Total	<0.0030	0.0069	0.1 ^a	0.1
Arsenic (As)-Total	0.0017	0.00192	0.005 ^a	0.005
Barium (Ba)-Total	0.0757	0.058	NS	NS
Beryllium (Be)-Total	<0.00010	<0.00010	100 ^b	NS
Boron (B)-Total	0.086	0.09	1.5 ^a	1.5
Cadmium (Cd)-Total	<0.0000050	0.0000155	0.00009 ^a	0.00033
Chromium (Cr)-Total	<0.00010	<0.00010	NS	NS
Cobalt (Co)-Total	<0.00010	<0.00010	0.05 ^a	0.0012
Copper (Cu)-Total	0.0292	<0.00050	0.0040 ^a	0.022
Iron (Fe)-Total	0.032	0.015	0.3 ^a	0.3
Lead (Pb)-Total	0.000201	<0.000050	0.007 ^a	0.007
Lithium (Li)-Total	0.0372	0.0404	2.5 ^b	NS
Manganese (Mn)-Total	0.557	0.0469	0.2 ^b	NS
Mercury (Hg)-Total	<0.0000050	<0.0000050	0.000026 ^a	NS
Molybdenum (Mo)-Total	0.00022	0.000186	0.073 ^a	0.073
Nickel (Ni)-Total	<0.00050	<0.00050	0.150 ^a	0.11
Selenium (Se)-Total	<0.000050	<0.000050	0.001 ^a	NS
Silver (Ag)-Total	<0.000010	<0.000010	0.00025 ^a	0.00025
Thallium (Tl)-Total	<0.000010	0.000011	0.0008 ^a	0.0008
Tin (Sn)-Total	<0.00010	<0.00010	NS	NS
Titanium (Ti)-Total	<0.00030	<0.00030	NS	NS
Uranium (U)-Total	0.000215	0.000191	0.01 ^b	0.015
Vanadium (V)-Total	<0.00050	<0.00050	0.1 ^b	NS
Zinc (Zn)-Total	0.0291	0.0056	0.007 ^a	0.03

1: CCME C Guidelines, de-minimis criteria for Protection of Aquatic Life and Agricultural Water Uses

2 - Environmental Quality Guidelines for Alberta Surface Waters 2018

a: CCME Canadian Environmental Quality Guidelines for water for the Protection of Aquatic Life

b: CCME Guidelines for Protection of Agricultural Water Uses (Irrigation and Livestock pathways included)

Table 6: Historical data of routine water chemistry in for Beaver Lake

Parameter	Year													
	1986	2005	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
pH	8.50	8.50	8.66	8.47	8.92	6.92	8.66	9.10	8.65	8.77	8.56	8.30	8.76	7.87
Secchi Depth (m)	2.90	2.60	4.17	3.56	1.85		1.53	1.88	1.50	1.88	1.50	2.40	2.50	2.70
Total Nitrogen (mg/L)	1.14	1.32	1.37	1.21	1.62	2.1	1.337	1.558	1.35	1.53	1.52	1.63	1.59	1.29
Total Phosphorus (mg/L)	0.033	0.145	0.047	0.024	0.065	0.050	0.041	0.039	0.040	0.034	0.036	0.040	0.029	0.031
Nitrate/Nitrite (mg/L)	0.0056	0.0018	0.0810	<0.071	<0.071		0.0	<0.022	<0.022	<0.022	<0.022	<0.022	<0.022	<0.022
Ammonia (mg/L)	0.0030	0.0331	0.0270	<0.050	<0.050	0.196	0.095	0.099	0.070	0.070	0.064	<0.050	<0.050	<0.05
Specific Conductivity (µS/cm)	409	462	508	500	529	600	554	491	559	493	757	526	887	729

Table 7: Historical data of total dissolved metals in Beaver Lake

Dissolved Metals	2018	2019	2020	2021	Criteria ¹	Criteria ²
	------(mg/L) -----					
Aluminum (Al)	0.0043	0.0094	<0.0675	0.0069	0.1 ^a	0.1
Arsenic (As)	0.00208	0.00207	0.00179	0.00192	0.005 ^a	0.005
Barium (Ba)	0.0602	0.0597	0.06445	0.058	NS	NS
Beryllium (Be)-Total	<0.00010	<0.00010	<0.00010	<0.00010	100 ^b	NS
Boron (B)	0.094	0.086	0.092	0.09	1.5 ^a	1.5
Cadmium (Cd)	0.0000055	<0.0000050	<0.0000050	0.0000155	0.00009 ^a	0.00033
Chromium (Cr)	0.00183	<0.00010	<0.00010	<0.00010	NS	NS
Cobalt (Co)-Total	<0.00010	<0.00010	<0.00010	<0.00010	0.05 ^a	0.0012
Copper (Cu)	0.016	<0.00050	<0.00050	<0.00050	0.0040 ^a	0.022
Iron (Fe)	<0.050	0.016	0.018	0.015	0.3 ^a	0.3
Lead (Pb)	0.000052	0.000060	<0.000635	<0.000050	0.007 ^a	0.007
Lithium (Li)-Total	0.0395	0.0403	0.0408	0.0404	2.5 ^b	NS
Manganese (Mn)	0.0998	0.0768	0.1522	0.0469	0.2 ^b	NS
Mercury (Hg)	<0.0000050	<0.0000050	<0.0000050	<0.0000050	0.000026 ^a	NS
Molybdenum (Mo)-Total	0.000177	0.000109	0.0001765	0.000186	0.073 ^a	0.073
Nickel (Ni)	0.00072	<0.00050	<0.00050	<0.00050	0.150 ^a	0.11
Selenium (Se)	<0.000050	<0.000050	<0.000050	<0.000050	0.001 ^a	NS
Silver (Ag)	<0.000010	<0.000010	<0.000010	<0.000010	0.00025 ^a	0.00025
Thallium (Tl)-Total	<0.000010	<0.000010	<0.000010	0.000011	0.0008 ^a	0.0008
Tin (Sn)-Total	0.0001	0.00010	<0.00010	<0.00010	0.0 ^a	NS
Titanium (Ti)-Total	<0.00030	0.00033	<0.00030	<0.00030	0.0 ^a	NS
Uranium (U)	0.000184	0.000162	0.0001835	0.000191	0.01 ^b	0.015
Vanadium (V)-Total	<0.00054	0.00113	<0.00050	<0.00050	0.1 ^b	NS
Zinc (Zn)	0.0092	<0.0030	0.0065	0.0056	0.007 ^a	0.03

1: CCME Canadian Environmental Quality Guidelines, de-minimis criteria for Protection of Aquatic Life and Protection of Agricultural Water

2 - Environmental Quality Guidelines for Alberta Surface Waters 2018

a: CCME Canadian Environmental Quality Guidelines for water for the Protection of Aquatic Life

b: CCME Guidelines for Protection of Agricultural Water Uses (Irrigation and Livestock pathways included)